


### Cuvierian tubules – where?

These unique structure are only found in some species of Aspidochirotida, more precisely in the genera:



## What are the Cuvierian tubules ?

They are intracoelomic caeca that generally occur in great number (up to 600) and attach to the basal part of the left respiratory tree.



## What is the function of the Cuvierian tubules ?

Cuvierian tubules are defensive organs.



Irritated, the holothuroids expel them through the anus

## What is the function of the Cuvierian tubules ?



The expelled tubules lengthen, become sticky and rapidly immobilize most organisms with which they come in contact!

Quiescent tubules are not sticky.

The elongated tubules do not adhere to mucus-cover surfaces, thus preventing them from sticking to the holothuroid body.



## How do Cuvierian tubules elongate?

- Expelled tubules are always in the process of elongation.
- Tubules elongate gradually from their basal extremity.
- Tubules elongation is irreversible and considerable (up to 20 times their original length!).
- Elongating tubules become progressively translucent while their diameter remains unchanged.




➔ **Elongation results from powerful entry of water into the tubule**

## How it works

The injection of water in the CT is as follow:

- ➔ Individuals completely contract their respiratory trees, expelling the water content in the cloaca.
- ➔ Part of the water expelled by the left respiratory tree enters the lumen of some tubules, causing their elongation.
- ➔ During contraction of the individuals, the hydrostatic pressure of the coelomic fluid increases and the opening of the anus causes cloacal rupture and tubules expulsion.

**Both tubules elongation and expulsion are basically caused by the contraction of the body wall which increases the pressure of the internal fluids.**



## Why do tubules lengthen?

From the inside to the outside, the tubules consist of:

a mesothelium lining the surface of the tubule that is exposed to the coelomic cavity.

a thick connective tissue layer



an inner epithelium surrounding the narrow lumen



## The inner epithelium



The lumen of the tubule is lined by a highly convoluted epithelium.





The basal part of epithelial cells consist of one to three elongated process penetrating deeply in the connective tissue.

The basal process contains piles of 5 to 25 spherules; the content of each spherule is released in the tubule lumen during elongation and could give the rigidity to an elongated tubule.

## The connective tissue

The connective tissue forms 90% of the tubule's thickness and encloses up to six imprecated collagen helices, each of them being parallel to the long axis of the tubule.




It also includes longitudinal and circular muscle fibers

## The mesothelium


The mesothelium is the tissue layer responsible for adhesion.

It is a pseudostratified epithelium made up of two superposed cell layers:




Granular cells are filled with densely packed granules enclosing the substance responsible of adhesion.

## Quiescent tubules - elongated tubules




During elongation, the structure of the Cuvierian tubule is modified



- The inner epithelium is dissociated,
- The helices of collagen fibers in the connective tissue layer are stretched,
- The protective outer layer of the mesothelium disintegrates
- The granular cell layer, now unfolded, becomes outermost on the tubule.

## Adhesion

Granular cells empty the contents of their granules only when the elongated tubule comes into contact with a surface, resulting in adhesion



Without contact

With contact

The adhesion of Cuvierian tubules is remarkable in several respects:

- ✓ adhesion occurs in seawater
- ✓ adhesion is achieved in a matter of seconds (less than 10 sec)
- ✓ adhesion remains active during several hours

Paradoxically, *the exact nature of the adhesive is still unknown!*


(More information in Flammang et al 2002)

## Maintaining the line of Defense

Expelled tubules autotomize at their attachment point on the left respiratory tree and are left behind as the holothuroid crawls away.

As only a portion of the tubules are emitted at one time, the total number may suffice for several responses.

After expulsion, the lost tubules are readily regenerated, making them a formidable defense mechanism.



(More information in VandenSpiegel et al 2000)

## Variation

All functional Cuvierian tubules (i.e. the expelled ones) investigated so far have almost the same structure.

The structure varies only in species where Cuvierian tubules are not expelled.

*Microthele*



*Actinopyga*




## The So-called Cuvierian organs in *Microthele*

### *Holothuria (Microthele)*

Some species of the subgenus *Microthele* possesses numerous branched tubules whose external surface shows alternative folds which are perpendicular to the long axis of the tubule.

The fine structure of the tubules closely resembles those of the *Holothuria* sp.. In contrast, basic behavioral differences occur as the tubules of *Microthele* sp. cannot elongated nor become sticky, or be expelled by individuals.


It is almost certain that these structure have no defensive role. Yet they are well developed organs that occur in great number in all observed individuals, but from the observation we made it is not possible to ascribe a possible function for these peculiar organs.



(More information in VandenSpiegel 1995)



## The So-called Cuvierian organs in *Actinopyga*



Cuvierian tubules

Cloaca

## The So-called Cuvierian organs in *Actinopyga*

Two main parts can be distinguished along an actinopygid Cuvierian tubule



The tubule branches


The tubule trunk



## The So-called Cuvierian organs in *Actinopyga*


### The tubule trunk

The trunk is a smooth tube with a narrow lumen, it cannot be elongated and has the overall tissue organization of the respiratory tree.




### The tubule branches


The tubule branches have no lumen and consists of a central rachis and many peripheral spherules.



The structure of the rachis resembles that of the tubule trunk.



The spherule itself is made of an enlarged spherical central core containing vesicular cells



## The So-called Cuvierian organs in *Actinopyga*

From the way actinopygid Cuvierian tubules attach and their overall histology they could be homologous to those of the other holothuroids but here again, it appears almost certain that these organs have no defensive role.

Additional information is required concerning the structure of the spherule's vesicular cells and their changes - for example during growth and/or during the annual cycle of the species - before a possible function can be ascribed to these peculiar organs

(More information in VandenSpiegel et al 1993)



**Thank you for your attention**